



**University of
Zurich**^{UZH}

**Zurich Open Repository and
Archive**

University of Zurich
University Library
Strickhofstrasse 39
CH-8057 Zurich
www.zora.uzh.ch

Year: 2012

Female philopatry and its social benefits among Bornean orangutans

van Noordwijk, M A ; Arora, N ; Willems, E P ; Dunkel, L P ; Amda, R N ; Mardianah, N ;
Ackermann, C ; Krützen, M ; van Schaik, C P

Abstract: Female philopatry in mammals is generally associated with ecological and sometimes social benefits, and often with dispersal by males. Previous studies on dispersal patterns of orangutans, largely non-gregarious Asian great apes, have yielded conflicting results. Based on 7 years of observational data and mitochondrial and nuclear DNA analyses on fecal samples of 41 adult Bornean orangutans (*Pongo pygmaeus wurmbii*) from the Tuanan population, we provide both genetic and behavioral evidence for male dispersal and female philopatry. Although maternally related adult female dyads showed similar home-range overlap as unrelated dyads, females spent much more time in association with known maternal relatives than with other females. While in association, offspring of maternally related females frequently engaged in social play, whereas mothers actively prevented this during encounters with unrelated mothers, suggesting that unrelated females may pose a threat to infants. Having trustworthy neighbors may therefore be a social benefit of philopatry that may be common among solitary mammals, thus reinforcing female philopatric tendencies in such species. The results also illustrate the diversity in dispersal patterns found within the great-ape lineage.

DOI: <https://doi.org/10.1007/s00265-012-1330-7>

Posted at the Zurich Open Repository and Archive, University of Zurich

ZORA URL: <https://doi.org/10.5167/uzh-61357>

Journal Article

Accepted Version

Originally published at:

van Noordwijk, M A; Arora, N; Willems, E P; Dunkel, L P; Amda, R N; Mardianah, N; Ackermann, C; Krützen, M; van Schaik, C P (2012). Female philopatry and its social benefits among Bornean orangutans. *Behavioral Ecology and Sociobiology*, 66(6):823-834.

DOI: <https://doi.org/10.1007/s00265-012-1330-7>

Female philopatry and its social benefits among Bornean orangutans

Authors:

Maria A. van Noordwijk¹, Natasha Arora¹, Erik P. Willems¹, Lynda P. Dunkel¹, Rahmalia N. Amda², Neneng Mardianah², Corinne Ackermann¹, Michael Krützen¹ and Carel P. van Schaik¹

Author Affiliation: ¹Anthropological Institute and Museum, University of Zurich, Switzerland; ²Fakultas Biologi, Universitas Nasional, Jakarta, Indonesia

Corresponding author: Maria van Noordwijk, Anthropological Institute and Museum, University of Zurich, Winterthurerstrasse 190, 8057 Zurich, Switzerland, Tel: +41-446355436, Fax: + 41-44 635 68 04
E-mail: vnoord@aim.uzh.ch

Key index words: male dispersal; female philopatry; pedigree; female-female association; range overlap; social play

Abstract

1 Female philopatry in mammals is generally associated with ecological and sometimes social benefits,
2 and often with dispersal by males. Previous studies on dispersal patterns of orangutans, largely non-
3 gregarious Asian great apes, have yielded conflicting results. Based on seven years of observational
4 data and mitochondrial and nuclear DNA analyses on fecal samples of 41 adult Bornean orangutans
5 (*Pongo pygmaeus wurmbii*) from the Tuanan population, we provide both genetic and behavioral
6 evidence for male dispersal and female philopatry. Although maternally related adult female dyads
7 showed similar home-range overlap as unrelated dyads, females spent much more time in association
8 with known maternal relatives than with other females. While in association, offspring of maternally
9 related females frequently engaged in social play, whereas mothers actively prevented this during
10 encounters with unrelated mothers, suggesting that unrelated females may pose a threat to infants.
11 Having trustworthy neighbors may therefore be a social benefit of philopatry that may be common
12 among solitary mammals, thus reinforcing female philopatric tendencies in such species. The results
13 also illustrate the diversity in dispersal patterns found within the great-ape lineage.

Introduction

In most mammals, females show philopatry, i.e. settle for life in or near the area in which they are born, whereas males disperse (Greenwood 1980; Waser & Jones 1983; Pusey and Packer 1987; Lawson Handley and Perrin 2007). Philopatry has ecological advantages in that it enables the individual to continue to live in a familiar habitat and maintain a familiar diet, without having to experience the costs in terms of time, travel and risk of finding a suitable area in which to settle. In addition, philopatry may enable individuals to maintain lifelong supportive social bonds with known relatives. In group-living primates, for instance, philopatric females tend to live in close association with their maternal relatives, who may provide mutual support in conflicts with less-related group members (e.g. Silk et al. 2009; 2010; see also Holekamp et al. 2011 for a similar system in spotted hyenas). Likewise, elephants living in fission-fusion female groups maintain supportive social relationships and show preferential associations with their matrilineal relatives in or near their natal range (De Villiers and Kok 1997; Archie et al. 2006). Thus, philopatry may bring social as well as ecological benefits. On the other hand, philopatry may limit access to unrelated and willing mates. Thus, in general, if members of one sex derive a clear advantage from being (more) philopatric, the other sex tends to derive a reproductive benefit from moving further away (Bengtsson 1978).

For species with strong ecological pressure on females to be more or less solitary, the social advantage of their philopatry is likely to be reduced, and therefore the tendency for female philopatry may be relaxed. Nonetheless, genetic and radio-telemetric research has confirmed that females in many non-gregarious mammal species settle in home ranges adjacent to, or overlapping with, their natal range and thus their female kin (e.g. tigers, Smith 1993; raccoons, Ratnayeke et al. 2002; bears, Støen et al. 2005; Moyer et al. 2006; woodrats, McEachern et al. 2007).

One possible explanation for the philopatric tendency of females in solitary mammals is that the ecological benefits on their own are strong enough to favor female philopatry, especially in species where males have much larger ranges than females and do not derive reproductive benefits from remaining philopatric. Where males derive strong benefits from philopatry, for example through coalitionary defense of a range or access to females, the less gregarious females are likely to disperse (e.g. chimpanzees, Langergraber et al. 2007; spider monkeys, Di Fiore et al. 2009). However, it is also possible that ‘solitary’ females in many species do accumulate social benefits, which accrue at rare but critical times in the life cycle. Indeed, there is evidence that, in some solitarily foraging species, females may be temporarily gregarious when they have dependent offspring. For example, females of the nocturnal mouse-lemur, *Microcebus murinus*, may share a nest hole for their young and even allo-nurse each other’s young. Only closely related females have been observed to share in this way,

thus both gaining foraging efficiency and maybe increased survival of their offspring through better thermoregulation and increased chances of adoption (Eberle and Kappeler 2006; cf. König 2006 for house mice). Some bats also associate at the roost preferentially with close maternal relatives during lactation, mutually gaining thermoregulatory benefits during a period of high energetic demands (Kerth et al. 2002). These observations raise the question whether social benefits are more common than usually assumed, which would strengthen a female philopatric tendency even in largely solitary species. Unfortunately, for most species it is unclear whether and how females interact differently with their related neighbors compared to unrelated ones, and thus whether they derive social benefits from philopatry at some stage during their lives.

Sumatran (*Pongo pygmaeus*) and Bornean (*P. abelli*) orangutans are the only ape species in which both males and females are habitually non-gregarious, like in many non-primate mammals. Especially the Bornean species shows a strong tendency towards solitary life (van Schaik 1999; van Noordwijk et al. 2009). This solitary lifestyle suggests that advantages for either sex of philopatry would be limited to the ecological advantage of familiar range and diet, unless unexpected social benefits are present. For this reason it is interesting to assess the degree to which females or males remain philopatric, and relate these tendencies to a detailed study of their social behavior.

Another reason for interest in orangutan philopatry patterns is that they are one of our closest living relatives. In chimpanzees, males are found to be the more gregarious sex and to be strongly philopatric, allowing them to form long-lasting bonds with their relatives, whereas females tend to disperse from their natal community (Langergraber et al. 2007). In another close relative, the gorilla, females as well as males usually disperse (e.g. Douadi et al. 2007), whereas males in Eastern gorillas may occasionally ‘inherit’ the groups in which they are born (Watts 2000; Bradley et al. 2007). Thus among the extant African apes at least female-biased dispersal seems to be the dominant pattern and has been proposed to be the tendency in the last common ancestor of humans and African apes (Ghiglieri 1987). Although various authors have assumed that female dispersal is a deeply rooted hominoid tendency (e.g. Foley and Lee 1991; Hrdy 2009; Chapais 2010), the ‘natural’ human dispersal system has remained a topic of debate (Hill et al. 2011). Therefore, establishing the natural dispersal pattern of the orangutan may provide a broader perspective to the reconstruction of the origins of the various dispersal patterns found within the hominoid lineage..

Long-term field observations on orangutan populations on both Sumatra and Borneo indicate that adult males have very large overlapping ranges, but rarely associate with each other (Galdikas 1985; van Schaik 1999; Utami Atmoko et al. 2009), and concentrate their associations with females largely to the periods in which these are potentially fertile. There is no evidence for male social bonds, as in chimpanzees, and thus male philopatry is

not expected. Females have smaller ranges than males (Singleton et al. 2009; Utami Atmoko et al. 2009), which can also largely overlap, and they spend most of their time accompanied only by their dependent (and sometimes one semi-dependent) offspring (van Noordwijk et al. 2009). For Sumatran orangutans, Singleton and van Schaik (2001; 2002) found clusters of females who associated more often with each other than expected based on their (extensive) range overlap. If females from the same cluster met, they were also more tolerant of close proximity than when females of different clusters encountered each other. These authors hypothesized that such female clusters were based on relatedness, thus implying female philopatry with some social benefits, but lacked the genetic data to verify this. On the other hand, Knott et al. (2008) emphasized for a Bornean population (Gunung Palung) that females show active avoidance of other females within their overlapping ranges, even though assumed relatives had more frequent encounters. In addition, they concluded that the outcome of a female-female encounter depended on its location relative to the females' respective core areas and thus that females defended their ranges. Thus, although female philopatry may be present in both orangutan species, social benefits appeared to be modest at best among Bornean orangutans.

Behavioral studies on maturing individuals have suggested a tendency towards female philopatry and male dispersal in orangutans of both islands (van Schaik and van Hooft 1996). However, the first studies that estimated genetic relatedness patterns among males and females in several populations have reached divergent conclusions, ranging from both sexes dispersing equally (Utami et al. 2002) to being equally philopatric (Goossens et al. 2006) to male-biased dispersal with female philopatry (Morrogh-Bernard et al. 2010). Nevertheless, recent population genetic analyses using both mitochondrial DNA and Y-chromosome markers indicate a spatial structuring congruent with male dispersal over large distances and very restricted female dispersal for both species (Arora et al. 2010; Nater et al. 2011; Nietlisbach 2009).

The aim of the present study was to use a combination of genetic, ranging and socio-behavioral data of individuals in an intensively studied population of orangutans (*Pongo pygmaeus wurmbii*) in Tuanan (Central Kalimantan, Indonesia) to assess for this population (i) whether females and/or males are philopatric or disperse, and (ii) whether philopatry was accompanied by social advantages. The genetic analyses focused on detecting mother – adult offspring dyads based on mitochondrial and nuclear DNA (for details see Arora et al. submitted). Given the stability of adult females' ranges (e.g. Wartmann et al. 2010 for the same population), the mother's range is the best estimate of any adult individual's natal range and thus maternal relatedness among adults is essential for documenting dispersal patterns.

Materials and methods

Study population

The Tuanan Orangutan Research Area (2° 09' South; 114° 26' East) is located in formerly selectively logged swamp forest on shallow peat (<2m) in the Mawas Conservation Area, Central Kalimantan, Indonesia. The orangutan population here has been studied since June 2003 and has approximately 4.5 individuals per km² (van Schaik et al. 2005). Individuals born before the start of the observations were assigned to age-sex categories based on prior experience (*cf.* Wich et al. 2004). The study area of c. 750 ha (gradually enlarged to > 1000 ha) encompassed several complete ranges of adult females, but ranges of all known males extended beyond this limited area. In this population, adult females spent on average less than 20 % of their time in association with another adult conspecific (van Noordwijk et al. 2009).

Observations in the field followed the standardized orangutan protocol (see www.aim.uzh.ch/orangutannetwork). Analyses on adult female ranging, association pattern, activities and social interactions are based on more than 16,650 hours of focal observation on 8 different mothers collected by a well-trained team of observers from July 2003 to July 2010. Most data were collected during nest-to-nest follows lasting a maximum of 10 consecutive days per month per focal female. During nest-to-nest follows, observation time was counted from the moment the focal female left her night nest in the morning until she rested in a (new) night nest in the evening (average active time per day for adult females was 10h 50min; n=1,330).

Adult female home ranges were found to be highly stable over time (Wartmann et al. 2010). Here we included data on adult females (who had had at least one offspring and were habituated to human observers) if in total at least 250 hours of focal observation and ranging data was collected for that female (thus excluding 5 known adult females ranging at the periphery of the study area). For each dyad of females only those data collected during the same years were used for a dyad-specific sample (resulting in different sample sizes per dyad – see supplemental table 1). All but one focal female had a dependent unweaned offspring (*i.e.* < 7 yr old) for at least part of the data collection period, and three focal females were accompanied at least part of the time by a young infant as well as a weaned offspring. In total the regular focal females had 5 female and 4 male immature offspring.

Sightings of all identified individuals were recorded for each month throughout the study period. The percentage of months an individual was present in the study area was based on focal follows as well as sightings during focal follows of other individuals or other research activities in the forest. Observational data on males were collected in the same way as for females. In total, over 10,500 hours of focal data was collected on those males sighted during at least 10% of the observation months. Data for the two morphs of adult males are presented separately, *i.e.* for flanged (with full secondary sexual characteristics, including cheek flanges) and

unflanged males (without such characteristics, but capable of siring offspring under natural conditions: Utami et al. 2002; Goossens et al. 2006). All young adult males are unflanged, but the age at which an individual male develops flanges is probably highly variable (Utami Atmoko et al. 2009).

Genetic Analyses

We evaluated the maternal relatedness of individuals at the study site through maternity analyses and inference of maternal siblings, using biparentally inherited microsatellite markers and a maternally transmitted mitochondrial DNA (mtDNA) marker. Fecal sample collection and storage was carried out using a standard genetic sampling protocol <http://www.aim.uzh.ch/orangutannetwork/GeneticSamplingProtocol.html>, followed by the generation of data on autosomal microsatellites and maternally inherited mitochondrial DNA (mtDNA) (details in Arora et al. 2010). We verified that the samples belonged to distinct individuals by genotyping them at six microsatellite markers, which had a combined non-exclusion probability of 1.36×10^{-5} and 8.90×10^{-3} for unrelated individuals and full siblings, respectively, as calculated in Cervus 3.0 (Kalinowski et al. 2007). This procedure identified 47 unique individuals. For 41 of these individuals we generated genotypes at an additional panel of 18 loci, totaling 24 autosomal microsatellite markers. For six males low DNA quantity and quality did not allow us to complete the genotyping for all loci. Thus, for these males, only the identification panel of six markers was used, which in combination with the use of mtDNA markers was sufficient to exclude all but one mother-son dyad (see Arora et al. subm).

To identify mother-offspring pairs, we carried out maternity analyses using the likelihood approach implemented in Cervus 3.0 (Kalinowski et al., 2007), using the strict 95% confidence level. All individuals were assessed as potential offspring, but only sexually mature females were incorporated as candidate mothers. To determine critical values of the log-likelihood score for a 95% confidence parentage assignment, we ran 10,000 simulations with the following parameters: a minimum of 6 loci typed, and our genotyping error rate of 0.112% (see Arora et al. subm.), as empirically determined through the “repeat-genotyping” and “unintentionally re-sampled individuals” approaches described by Hoffman and Amos (2005). The proportion of candidate mothers sampled was difficult to estimate from field data. It has been shown that this parameter may have a substantial influence on the statistical significance of the parentage assignments (Krützen et al. 2004). Thus, we tested several conservative values (0.05, 0.08 and 0.10) and corroborated that the results were robust. In all cases, mother offspring dyads detected using microsatellite data, also shared their mtDNA haplotype. In addition, all known mother-unweaned offspring dyads, with samples available for both (N=8), were confirmed to have a genetic mother-offspring relationship according to our procedures. Individuals sharing a mother were inferred to

be maternal siblings. A limitation to this approach is that individuals that are maternally related may not be detected due to the absence of (a sample of) a shared mother. However, this bias is not expected to differ for females compared to males. For all further analyses, maternally related dyads were defined as either a mother-offspring pair, or two individuals sharing a mother.

Ranging behavior

The ranging behavior of 8 focal females was investigated at the level of both home range and core range areas to obtain estimates of proportional dyadic overlap. Using locational data collected at 30-minute intervals (see criteria for inclusion above), the respective ranging areas of all females were calculated on a dyad-specific basis. We delineated home ranges by 95% volume isopleths (Andersen 1982) on utilization distributions obtained from fixed Gaussian kernel density estimation (using BCV to estimate the kernel's bandwidth; Worton 1989), whereas core areas were defined by 50% volume isopleths. Areas of overlap were subsequently calculated and divided by the dyad-specific home range area of each focal female. This resulted in an asymmetric matrix for both proportional home range and core range overlap of 52 of the 58 possible dyads (insufficient concurrent ranging data were available for 6 dyads). All analyses on ranging behavior were conducted using the HRT plug-in (Rodgers et al. 2007) and Spatial Analyst extension for ArcGIS 9.3 (ESRI 2008).

Association and social interaction analyses

Whenever two individuals approached to within 50m this was considered an encounter. Encounter rates were based on the number of new approaches within 50m occurring during a focal follow starting at the morning nest. If a dyad had spent the night within 50m of each other, the (continuing) association the next morning was not counted as a new encounter. The percentage of time in association (< 50 m distance) was based on total active time from the focal female's perspective. The relationship between time in association and range overlap between two females was based only on those years for which ranges for both could be calculated. When individuals were simultaneously feeding in the same food source <10m apart they were said to show 'feeding tolerance'. An agonistic interaction was defined as one in which one individual shows clear aggressive acts such as slapping, grabbing, biting or a fast chase and/or the other shows obvious avoidance or submissive behavior such as fleeing fast (through canopy or over the ground) or screaming. In the analyses, social play among offspring of different mothers could include both unweaned and weaned offspring as long as these were in permanent association with the mother.

Unfortunately, as in previous studies on wild orangutans, the small number of mother-offspring dyads did not allow for an analysis of the effect of offspring sex (van Noordwijk et al. 2009). However, so far, no striking differences between female and male dependent offspring in time budgets or social interest have become apparent.

Statistical analyses

Given that dyadic data are inherently non-independent, permutation versions of standard statistical techniques were employed in which significance of test-statistics was assessed by 10'000 randomizations. To test whether the degree of overlap in ranging areas (both at the home and core range level) differed between related and unrelated female dyads, permutation unpaired t-tests were conducted (Legendre and Legendre 1998). Differences in the duration of association among dyads were compared in a permutation one-way ANOVA-test (Legendre and Legendre 1998). Post-hoc pairwise comparisons (Tukey's Honestly Significant Difference tests) subsequently revealed which categories of dyad were significantly different from each other. Potential associations between overlap in ranging areas and encounter frequency and association time were considered separate for unrelated and related female dyads by computing Pearson permutation correlation tests. Differences in the encounter frequencies and duration of association between unrelated and related female dyads were investigated in more detail by a permutation unpaired t-test.

Results

Genetic analyses

In total, 10 different mtDNA haplotypes were found for the individuals sampled in Tuanan (see Arora et al. subm. for details). All females (including all additional adolescent and nulliparous ones) had one of only 3 haplotypes, whereas for the males 8 different ones were found. Only one haplotype was shared by males and females. This distribution of haplotypes was significantly different for females and males (Arora et al. subm.).

Pedigree analysis based on nuclear and mtDNA indicated the presence of one mother with 3 adult daughters (for which maternal sibship was thus inferred) and one mother-adult daughter dyad, all with ranges mostly inside the study area. In addition, 2 independently ranging adolescent females could be matched with their mothers ranging in the periphery of the study area. In contrast among the 28 males, only 1 young one (estimated 10-15 years old) could be matched with his peripherally ranging mother. Thus, both the much greater concentration of females in a few haplotypes and diversity of haplotypes of the males and the sex difference in

the presence of dyads of maternally closely related adults in the study area are consistent with greater female philopatry and male dispersal (see also Arora et al. subm).

In the following sections we refer to the 4 genetically detected mother-adult daughter dyads as well as the 3 female-female sibling dyads (based on sharing the same mother) as ‘related dyads’ and all other female-female dyads as ‘unrelated dyads’.

Spatial analyses

Male ranges and overlap

None of the known adult males had his complete range within the study area. In addition, most of the known males were seen throughout this area. Thus, male ranges were estimated to be considerably larger than even the expanded study area of 1000 ha. In addition, all known males were regularly observed to leave the study area. Only one (flanged) male was sighted during c. 50% of the observation months, (compared to 4 females during >75% of the months), whereas 13 additional males (7 flanged; 5 unflanged) were sighted during at least 10% of the months (the other 14 genetically identified males [11 flanged and 4 unflanged] were sighted less often). These observations suggest major home range overlap among the many adult males sighted in the area.

Female ranges and overlap

Some females could never be followed for more than a few consecutive days before they left the study area, whereas others never travelled outside of the study area during follows. In the analyses of home range overlap we only included those females for whom we were confident to have adequate data to calculate their 95% and 50% use area, based on 1-8 years of data per dyad. Overlap was always calculated from the perspective of the focal female (resulting in 2 different values per dyad). The average home range size for females with ranges inside the study area was 327.5 ± 124.7 ha, with a Core Area (50% use) of on average 84.0 ± 27.6 ha.

All known maternally related female dyads had overlapping home ranges (Fig. 1; average HR overlap per dyad: 57.32%, N=12), as well as overlapping core areas (average 15.79% (Fig 2a). Within the study area, non-related female dyads (N= 40) had an average overlap of 36.90% and core range overlap of 6.85%, which is significantly less (home range overlap: Permutation t-test $t_{(50)}=2.51$, $P=0.015$, $n_{perm}=10'000$; Core range overlap: Permutation t-test $t_{(50)}=2.25$, $P=0.027$, $n_{perm}=10'000$). However, more than half of the unrelated dyads (N= 23) exhibited a degree of overlap equal to or greater than the minimum observed overlap between related dyads (32.1%), and when comparing these (average 54.04% HR overlap and 10.76% CR overlap) to related

dyads, differences were no longer significant; Permutation t-test for HR overlap: $t_{(33)}=0.466$, $P=0.65$; CR overlap $t_{(33)}=1.029$, $P=0.32$). Thus matrilineal clusters of females have overlapping ranges, but home ranges and core areas may be shared with unrelated adult females as well, and to a similar extent.

Social relationships

Male-male social relationships

During 7 years of focal sampling on adult males, male-male encounters and associations were rare. Flanged males spent on average 0.11 % of their focal time in association with another flanged male, flanged with unflanged males on average 0.25%, and unflanged with unflanged males 3.12%, but only 1.0% when they were not also in association with an adult female. Associations among male dyads of all combinations lasted significantly shorter than those among closely related females, but were not different from those of unrelated females (permutation one-way ANOVA: $F=20.70$, $P=0.0001$; post-hoc pairwise comparisons -Tukey's HSD- reported in Supplementary Table 2).

Only unflanged males occasionally showed feeding tolerance towards other unflanged males (at least briefly during 26% of unflanged male associations, or 8% of all male-male associations). Social play among 2 unflanged males was observed during 5 associations – (four times involving the same male with 3 different partners) and except for one event such play lasted only for a few minutes. No grooming, or coalitions against another individual were ever observed among males and only 1 association between unflanged males was continued the next morning (see also Supplementary Fig. 1). Thus, we found no evidence for any special relationships, social bonds or coordinated ranging behavior among adult males in this population.

Female social relationships

Female encounters and associations

The frequency of encounters amongst female dyads was not related to the percentage of home range overlap (permutation correlation test: related dyads: $r_{\text{Pearson}}=-0.22$, $N=12$, $P_{\text{perm}(n=10'000)}=0.49$; unrelated dyads: $r_{\text{Pearson}}=0.19$, $N=40$, $P_{\text{perm}(n=10'000)}=0.24$). Even degree of core range overlap was not significantly correlated with encounter frequency among related dyads ($r_{\text{Pearson}}=-0.16$, $N=12$, $P_{\text{perm}(n=10'000)}=0.62$), whereas it was positively correlated for unrelated dyads ($r_{\text{Pearson}}=0.41$, $N=40$, $P_{\text{perm}(n=10'000)}=0.019$). However, if we only consider those unrelated dyads with a total home range overlap equal to or greater than the minimum observed amongst related females, there was only a trend ($r=0.34$, $N=23$, $P_{\text{perm}(n=10'000)}=0.09$).

Overall, females encountered their close maternal relatives more often than other females with similar degree of home range overlap, i.e. those unrelated dyads with an overlap of more than the minimum among related dyads of 32.1% (Permutation unpaired t-test: $t=6.49$, $N_1=12$, $N_2=23$, $P=0.0001$). This distinction between relatives and other females is also reflected in the percentage of time focal females spent in association with other adult females (Fig 2b; $t=5.59$, $N_1=12$, $N_2=23$, $P=0.0001$). Additional analyses of encounter rate and association duration divided by percentage home range and core range overlap respectively, yielded also very significant differences between related and unrelated dyads (Fig 2c).

Once associations occurred, their duration among related females was also longer than among non-related females (Supplementary Table 2). In addition, none of the 37 observed associations between unrelated females lasted for more than 3 hours and none was maintained overnight, whereas 96 of 194 (49.7%) associations of related females lasted more than 3 hours, and in 54 out of 194 (27.8%) a night nest was made within 50m of each other. This bias in both encounter frequency and association time strongly suggests that spending time in association is an active choice and not a by-product of range overlap.

Social interactions during associations

Agonism. Even if we restrict analyses to the relatively short associations (< 3h) observed among unrelated females, associations among non-kin dyads differed from those among kin-dyads (Fig 3), with a higher frequency of severe agonistic interactions (i.e. with active chase and or physical fight including hitting or biting: in 8 out of 37 associations vs. 5 out of 98 Fisher exact $P=0.008$). In addition, agonistic interactions among non-relatives always (8/8) led to immediate termination of the association and only in 2/5 of the cases among related dyads. (All severe agonism among relatives was between the same two half-sisters).

Feeding tolerance. During associations females hardly ever engaged in positive social activities (only once brief grooming in a mother-adult daughter dyad), but sometimes showed tolerance by feeding within 10 m of each other in the same food patch. Such 'feeding tolerance' was never seen during associations of non-related dyads, but in 32 of 98 (32.7 %) of those of maternal relatives ($\chi^2=14.08$; $P<0.001$; Fig 3).

Social play. Despite the females' lack of affiliative interactions during most associations, their respective offspring frequently engaged in social play (mostly arboreal play-wrestling). Related mothers rarely terminated, and sometimes even actively enabled play among peers while protecting very young infants, and mothers, especially of infants <3 yr of age, sometimes even actively participated in such play. However, mothers intervened and effectively prevented contact among non-kin peers on several occasions, by either retrieving their own offspring or chasing away the other immature if it was accompanied by its mother. As a result, social play

between unweaned immatures was only seen (briefly) during 3 (of 37) associations between non-related female dyads, whereas play was observed during 40 (of 98) of the associations between relatives ($\chi^2=11.77$; $P<0.001$; comparing only associations lasting < 3 h to decrease bias in favor of related dyads; Fig 3). Nevertheless, when weaned immatures ranging independently from their own mother (>50 m away) visited a non-related mother-offspring pair, they were ‘allowed’ to play with dependent offspring (observed during at least 6 different focal follows), even for a prolonged period (average play duration 60 min.).

In summary, encounters between unrelated females rarely resulted in long-lasting association, and never included simultaneous feeding in the same food patch. Agonistic termination of such associations was also more likely than of associations among related females. In contrast, the close kin associations are characterized by tolerance during feeding and tolerance and accommodation of play interactions among their offspring.

Female cluster size and infant playing time

The Tuanan study area included the ranges of one matrilineal kin-group with a mother and her 3 adult daughters and another mother-adult daughter dyad ranging at the edge of the study area (insufficient data on 1 of these 2 females). In addition, data were collected on 3 adult females ranging mostly (but not completely) in the study area, and for these females no living adult maternal relatives were known. Even though these females may have had relatives living outside the study area, those were never encountered during follows. Thus in the study area, immatures had access to kin clusters of different sizes, and therefore they were likely to grow up with different opportunities for social interactions with peers. Indeed, in this study females and their offspring of the larger kin cluster with 4 adult females spent more time in total in association with other adult females than those with only one or no known adult maternal relatives (average 6.15 ± 3.07 % vs. 0.34 ± 0.08 %; Mann Whitney $U=0$, $N_1=4$, $N_2=4$, $P < 0.05$ two-tailed; whereas large and small kin cluster females did not differ in % time in association with unrelated females: $U=10$, $N_1=4$, $N_2=4$, NS). Even though this comparison is based on only one large maternal cluster vs. several females from small clusters, these data suggest a matriline size effect on social opportunities for maturing offspring.

The time difference in time spent in association with peers (and their mothers) was reflected in the time budgets for 1-5 year old dependent immatures, indicating that immatures growing up in the large kin cluster consistently spent more time in social play than immatures growing up in a small kin cluster (average 1.18 ± 0.92 % $N=11$ yearly values of at least 150hr of focal data for the large cluster vs. 0.06 ± 0.06 % $N=5$ for the small; Mann Whitney $U=4$, $N_1=11$, $N_2=5$, $P < 0.01$ two-tailed). Despite the small sample sizes, these data strongly suggest that maternal cluster size affects the amount of social play with peers an immature can achieve.

Discussion

Dispersal and philopatry

We used observational and genetic data from a single study area to investigate the relatedness patterns among the resident adults in a non-gregarious great ape. Genetic analyses focused on close maternal relatedness, since female ranges were very stable and the maternal range is therefore the best predictor of an adult's natal range. The combination of ranging data and the genealogical reconstruction of all adults sighted in the study area indicated that whereas 4 parous (plus at least 2 nulliparous) females lived in overlapping ranges with their surviving mother, we found only one young male being maternally related to a known female at the periphery of the study area. Thus, our results strongly suggested female philopatry (*sensu* Waser & Jones 1983) and male dispersal and are therefore consistent with other recent genetic studies (Arora et al. 2010; Morrogh-Bernard et al. 2010; Nater et al. 2011).

From the Tuanan results, we could also estimate the age at which males disperse from their natal area. The one male with a female relative in the study area was still young (estimated <15 yr old) and never seen in the same area or in association with his mother. At least 4 unflanged males had a 'nonlocal haplotype', e.g. different from all of the sampled parous females (see Arora et al. *subm.*). These 4 males were frequently present (10-50% of the months) in the study area, indicating they had moved away from their natal area while they were still unflanged. We therefore conclude that males tend to disperse from their natal range as adolescents. Morrogh-Bernard et al. (2010) concluded that males do not disperse before growing flanges. However, their conclusion was based on a small sample, and is not easily reconciled with the fact that unflanged males are known to sire offspring (Utami et al. 2002; Goossens et al. 2006). We therefore suspect that our conclusion holds more generally.

Recent genetic analyses of the Y-chromosome suggest that males may disperse remarkably long distances away from their natal areas (Nietlisbach 2009). However, here we found that male orangutans not only disperse spatially (away from natal area), they also do so socially (away from known relatives) (*cf.* Isbell and van Vuuren 1997). First, among the males in the study area, no close maternal male kin could be identified. Second, male-male associations, even among unflanged males, were of significantly shorter duration than associations between related females, and no consistent associates could be detected, suggesting the absence of the social bonds expected if there was parallel dispersal (see also Supplementary Fig. 1). Females, in contrast, appear to stay in the familiar area and also near their familiar female relatives, although we would need to sample a larger area to assess whether all females manage to do so. Overall, therefore, the genetic results

strongly support behavioral evidence for female philopatry and continuing association with relatives in Bornean orangutans, and the opposite pattern in males.

Female relationships and the social benefits of philopatry

It is possible that maturing individuals with a greater number of tolerant role models would have acquired a larger number of learned skills by the time they are adult. Although infants peer largely toward the activities of their own mothers they also do so occasionally with other cluster members (Jaeggi et al. 2010), but never with females of other clusters. However, given the small sample sizes, the fitness benefit is hard to quantify. Future work will attempt to estimate these benefits.

In this study, the most conspicuous behavior during associations was social play. Even though related female dyads seemed to choose to approach and spend time in proximity, adults tended to watch each other initially and only gradually approach to within 5m. Females' dependent offspring, however, tended to approach each other quickly and start social play within the first minutes of the start of the association. Successful maternal intervention, by chasing the other immature or retrieving her own offspring, was seen on several occasions when unrelated females were in brief association and their infants approached each other. The few observations of social play between unrelated immatures almost all happened in the absence (> 50m away) of one of the mothers, or when one of the mothers was distracted by a consortship with a male. Lack of maternal proximity and attention only occurred for immatures of at least 5 years old. This pattern suggests that the lack of play among peers born in different maternal clusters is due to the mothers' reluctance to allow their offspring to interact and not to a lack of interest on the part of the immatures.

Opportunities for social play among peers are rare for this sample of wild Bornean orangutans. Even though some mothers occasionally actively engaged in social play with their young offspring, they certainly did not do this every day. An older sibling, if present, is at least 7 years older and thus much larger and, most importantly, no longer in frequent association by the time the infant is 2-3 years old and ready to move more than a few meters away from the mother (van Noordwijk et al. 2009). However, when related mother-offspring pairs were in association, immatures frequently played and seemed to forego not only solitary play (which accounts for 20-50% of a 1-4 yr olds' average time budget), but also reduced their time feeding and resting (sometimes by >10%) (unpublished data). Even before weaning, social play in this population did not exceed 4 % of the average yearly time budget, whereas it drops to less than 1 % after weaning (unpublished data). Nevertheless, immatures seem to take advantage of every opportunity they can get to engage in social play with peers, whereas during their associations, adult females rarely engage in social exchanges (such as grooming or

food sharing) and merely tolerate proximity (or not). Thus association among parous females seems to be in the interest of the offspring more than of the mothers themselves.

Why would it be beneficial for mothers to provide opportunities for social interactions with peers?

Social play, especially during development, is seen in all primates and indeed most mammals and birds as well as other vertebrates (Fagen 1981; Graham and Burghardt 2010). The major functions of play are thought to be facilitation of motor development (e.g. Byers and Walker 1995; Nunes et al. 2004), and brain development (e.g. Lewis and Barton 2006; Pellis et al. 2010), and preparing the individual to respond to unexpected events (Špinka et al. 2001). Yet, the fitness consequences of a lack of social play under natural conditions are still little known and hard to measure, although one study on Alaskan brown bears, (Fagen and Fagen 2009) reported a positive correlation between social play and survival to independence in Alaskan brown bears, irrespective of food availability and maternal condition. However, despite several plausible hypotheses about the benefits of social play, it remains to be determined whether the difference we found in this study between a little bit of play (on average c. 1 hour per week) for the ‘large-kin-cluster’ immatures and virtually nothing (a few minutes per week) for the ‘small-kin cluster’ immatures, could affect the orangutan immatures’ (social) development and, therefore, be biologically meaningful.

Assuming that social play is important for the development of their offspring, why would mothers not tolerate play among non-relatives, or only when the other mother is absent? When an independent immature ‘visits’ a mother offspring pair, the mother, being larger, can easily intervene in the social interaction whenever her offspring signals distress or is risking injury by falling out of the trees. However, if another adult female is also present, she may be less able to quickly rescue her offspring. This might explain why females tend to tolerate the offspring’s interactions with well-known maternal relatives, who benefit to some extent from the well-being of both immatures, but are wary or antagonistic towards unrelated females with whom they don’t share a common interest. Another explanation may be that it is generally costly for females to spend time in association (potential scramble competition) and when they do this for social reasons (or their offspring’s social benefit) prefer a familiar association partner. Familiar partners are likely to impose the lowest physiological cost (lowest stress response) by being predictable and having some shared interest.

Among East African chimpanzee females, who can be almost as non-gregarious as orangutans, severe female aggression has been documented, resulting in the death of a competitor’s offspring (Goodall 1986; Pusey et al. 1997; Townsend et al. 2007). Even though orangutans have never been seen to form coalitions in the wild, as in the documented cases of female infanticide in chimpanzees, females have been seen to fight (including biting) with each other on rare occasions. Thus the potential for serious harm to unprotected young immatures is

clearly present. In general, mistrust between unrelated females is expected because of the potential harm competing females could do to each other's offspring (cf. offspring-defense hypothesis: Wolff & Peterson 1998), whereas tolerance is found to be higher among related neighbors (Waser & Jones 1983; Kitchen et al. 2005). It is noteworthy that orangutan females without offspring (nulliparous and females with only older weaned offspring) were more likely to range over a larger area, but retreated into their established range after the birth of their next offspring (van Woerden and Pettersson 2007, and unpublished data). A likely benefit of the more restricted ranging of mothers with young offspring may be that within their smaller range the risk of encounters with strangers is smaller. We have shown here that females respond very differently to encounters with maternally related females vs. maternally not related females. This suggest a potential risk posed by unrelated females, even though this may be hard to document.

Dispersal and female relationships in the great ape lineage

This study showed that females of the least gregarious extant great ape actually do maintain social relationships with their philopatric maternal female kin and so seem to enable the development of social bonds among their offspring. In this respect, orangutans are very similar to many other primates living in matrilineal groups, but different from the African great apes which have a tendency for female dispersal with limited opportunity for female bonding. Thus, these results illustrate the variability in philopatric tendencies among all great apes from strongly male-biased to strongly female-biased dispersal.

Perhaps more importantly, the results underscore the fact that orangutan females, in spite of their philopatric tendency, may occasionally live without any nearby adult female relatives. However, the long-term fitness consequences for their offspring remain to be examined. Among the other great apes, females seem to be even more flexible. For example, although chimpanzee females are sometimes able to stay in their natal community and then maintain matrilineal bonds (Goodall 1986; Pusey et al. 1997), they can also form supportive dyadic relationships with particular unrelated females after dispersal (Wakefield 2008; Langergraber et al. 2009; Wild 2010). Bonobo females also form close bonds with unrelated females (Hashimoto et al. 1996) and gorilla females are sometimes able to maintain social bonds with relatives as well as with unrelated females (Watts 1994; Bradley et al. 2007). This ability to form and maintain bonds, has freed females from the necessity to be strictly philopatric.

Similar flexibility and independence from philopatry for the formation of intrasexual bonds is also seen among human females. Hill et al. (2011) recently showed that among hunter-gatherers, an individual's or even a female-male pair's site of residence may vary throughout the lifetime. This unusually flexible pattern of

dispersal enables a young mother to live near maternal kin and receive their support when she most needs it, but also enables males to co-reside with male kin and form coalitions at other times in their life. As data on great apes accumulate, a similar flexibility in philopatry and dispersal is becoming apparent.

Acknowledgements

We are grateful to the Indonesian Institute of Science (LIPI), and the Indonesian State Ministry for Research and Technology (RisTek), the Director General Departemen Kehutanan (PHKA), Departamen Dalam Negri, the local government in Central Kalimantan, the BKSDA Palangkaraya, the Bornean Orangutan Survival Foundation (BOSF) and MAWAS in Palangkaraya for their permission and support to conduct this research. We also thank the Fakultas Biologi Universitas Nasional (UNAS) in Jakarta for their collaboration and support for the Tuanan project and in particular drs. Tatang Mitra Setia and dr. Suci Utami Atmoko, and we gratefully acknowledge the collaboration of dr. Joko Pamungkas of PSSP-IPB with permit arrangements for the genetic samples. We are indebted to field team and in particular Pak Rahmatd and Pak Yandi for their help with collecting data and samples in the field. We thank Claude Rosselet and Pascal Marty for their work on the ArcGis database, and Alex Nater, Elvira Schneider and Moritz Fischer for their contributions to the genetic analyses. We thank the reviewers and David Watts for their constructive comments on the manuscript. Samples were exported from Indonesia to Switzerland under permits 07279/IV/SATS-LN/2009, 00961/IV/SATS-LN/2007 from the Convention on International Trade in Endangered Species of Wild Fauna and Flora. For major financial support we thank the University of Zurich, and the A.H. Schultz Stiftung.

Ethical standards

Behavioral observations comply with the Indonesian laws. The authors declare no conflict of interest.

References

- Anderson DJ (1982) The home range: a new nonparametric-estimation technique. *Ecology* 63:103-112
- Archie EA, Moss CJ, Alberts SC (2006) The ties that bind: genetic relatedness predicts the fission and fusion of social groups in wild African elephants. *Proc Roy Soc B* 273:513-522
- Arora N, Nater A, van Schaik CP, Willems EP, van Noordwijk MA, Goossens B, Morf N, Bastian M, Knott C, Morrogh-Bernard H, Kuze N, Kanamori T, Pamungkas J, Perwitasari-Farajallah D, Verschoor E, Warren K, Krützen M (2010) Effects of Pleistocene glaciations and rivers on the population structure of Bornean orangutans (*Pongo pygmaeus*). *Proc Natl Acad Sci* 107: 21376-21381
- Arora N, van Noordwijk MA, Ackermann C, Willems EP, A. Nater A, Greminger M, Nietlisbach P, Dunkel LP, Utami Atmoko SS, Perwitsari-Farajallah D, van Schaik CP, Krützen M (submitted) Parentage-

511 based pedigree reconstruction reveals female matrilineal clusters and male-biased dispersal in non-
 512 gregarious Asian great apes, the Bornean orangutans (*Pongo pygmaeus*).
 513 Bengtsson BO (1978) Avoiding inbreeding - at what cost? J Theor Biol 73:439-444
 514 Bradley BJ, Doran-Sheehy DM, Vigilant L (2007) Potential for female kin associations in wild western gorillas
 515 despite female dispersal. Proc Roy Soc B 274: 2179-2185
 516 Byers JA, Walker C (1995) Refining the motor training hypothesis for the evolution of play. Am Nat 146:25-40
 517 Chapais B (2010) The deep structure of human society: primate origins and evolution. In: Kappeler PM, Silk JB,
 518 (eds) Mind the gap. Tracing the origins of human universals. Springer Verlag, Berlin, pp 19-51
 519 De Villiers PA, Kok OB (1997) Home range, association and related aspects of elephants in the eastern
 520 Transvaal Lowveld. Afr J Ecol 35:224-236
 521 Di Fiore A, Link A, Schmitt CA, Spehar SN (2009) Dispersal patterns in sympatric woolly and spider monkeys:
 522 integrating molecular and observational data. Behaviour 146:437-470
 523 Douadi MI, Gatti s, Levrero F, Duhamel G, Bermejo M, Vallet D, Menard N, Petit EJ (2007) Sex- biased
 524 dispersal in western gorilla (*Gorilla gorilla gorilla*)Molecular Ecology 16: 2247–2259 doi: 10.1111/j.1365-
 525 294X.2007.03286.x
 526 Eberle M, Kappeler PM (2006) Family insurance: kin selection and cooperative breeding in a solitary primate
 527 (*Microcebus murinus*). BehavEcol and Sociobiol 60:582-588
 528 ESRI (2008) ArcGis 9.3. Environmental Systems Research Institute Inc, Redlands, CA
 529 Fagen J (1981) Animal play behavior. Oxford University Press, New York
 530 Fagen R, Fagen J (2009) Play behaviour and multi-year juvenile survival in free-ranging brown bears, *Ursus*
 531 *arctos*. Evol Ecol Res 11:1053–1067
 532 Foley RA, Lee PC (1991) Ecology and energetics of encephalization in hominid evolution Phil Trans Roy Soc B
 533 334:223-232
 534 Galdikas, BMF (1985) Orangutan sociality at Tanjung Puting. Am J Primatol 9: 101-119
 535 Ghiglieri MP (1987) Sociobiology of the great apes and the hominid ancestor. J Hum Evol 16: 319-357
 536 Goodall J (1986) The chimpanzees of Gombe. Harvard University Press, Cambridge, MA
 537 Goossens B, Setchell JM, James SS, Funk SM, Chikli L, Abulani A, Ancrenaz M, Lackman-Ancrenaz I, Bruford
 538 MW (2006) Philopatry and reproductive success In Bornean orangutans (*Pongo pygmaeus*). Mol Ecol
 539 15:2577-2588
 540 Graham KL, Burghardt GM (2010) Current perspectives on the biology study of play: signs of progress. Q Rev
 541 Biol 84:1-24
 542 Greenwood PJ (1980) Mating systems, philopatry and dispersal in birds and mammals. Anim Behav 28:1140-
 543 1162
 544 Hashimoto C, Takenaka O, Furuichi T (1996) Matrilineal kin relationship and social behavior of wild bonobos
 545 (*Pan paniscus*): Sequencing the D-loop region of mitochondrial DNA. Primates 37:305-318
 546 Hill KR, Walker RS, Božičević M, Eder J, Headland T, Hewlett B, Hurtado AM, Marlowe F, Wiessner P, Wood
 547 B. (2011) Co-residence patterns in hunter-gatherer societies show unique human social structure.
 548 Science 311:1268-1289
 549 Hoffman JI, Amos W (2005) Microsatellite genotyping errors: detection approaches, common sources and
 550 consequences for paternal exclusion. Mol Ecol 14, 599-612
 551 Hrdy SB(2009) Mothers and Others: the Evolutionary Origins of Mutual Understanding. Cambridge MA:

552 Harvard University Press.
 553 Isbell LA, Van Vuren D (1996). Differential costs of locational and social dispersal and their consequences for
 554 female group-living primates. *Behaviour* 133:1-36
 555 Jaeggi AV, Dunkel LP, van Noordwijk MA, Wich SA, Sura AAL, van Schaik, CP (2010) Social learning of diet
 556 and foraging skills by wild immature Bornean orangutans: implication for culture. *Am J Primatol* 72:
 557 62-71
 558 Kalinowski ST, Taper ML, Marshall TC (2007) Revising how the computer program CERVUS accommodates
 559 genotyping error increases success in paternity assignment. *Mol Ecol* 16:1099-1106
 560 Kerth G, Safi K, König B (2002) Mean colony relatedness is a poor predictor of colony structure and female
 561 philopatry in the communally breeding Bechstein's bat (*Myotis bechsteinii*). *Behav Ecol Sociobiol*
 562 52:203-210
 563 Kitchen AM, Gese EM, Waits LP, Karki SM, Schauster ER (2005) Genetic and spatial structure within a swift
 564 fox population. *J Anim Ecol* 74: 1173-1181
 565 Knott CD, Beaudrot L, Snaith T, White S, Tschauner H, Planasky G (2008) Female-female competition in
 566 Bornean orangutans. *Int J Primatol* 29:975-997
 567 König B (2006) Non-offspring nursing in mammals: General implications from a case study on house mice. In:
 568 Kappeler PM van Schaik CP (eds) *Cooperation in Primates and Humans Mechanisms and Evolution*.
 569 Springer-Verlag, Berlin, pp 191-205
 570 Krützen M, Barre LM, Connor RC, Mann J, Sherwin WB (2004) 'O father: where art thou?'--Paternity
 571 assessment in an open fission-fusion society of wild bottlenose dolphins (*Tursiops* sp.) in Shark Bay,
 572 Western Australia. *Mol Ecol* 13, 1975-1990
 573 Langergraber K, Mitani J, Vigilant L (2009) Kinship and social bonds in female chimpanzees (*Pan troglodytes*).
 574 *Am J Primatol* 71:840–851
 575 Langergraber KE, Mitani JC, Vigilant L (2007) The limited impact of kinship on cooperation in wild
 576 chimpanzees. *Proc Natl Acad Sci* 104:7786-7790
 577 Lawson Handley LJ, Perrin N (2007) Advances in our understanding of mammalian sex-biased dispersal. *Mol*
 578 *Ecol* 16:1559-1578
 579 Legendre P, Legendre L (1998) *Numerical Ecology*. 2nd edition. Elsevier, Amsterdam
 580 Lewis KP, Barton RA (2006) Amygdala size and hypothalamus size predict social play frequency in nonhuman
 581 primates: A comparative analysis using independent contrasts. *J Comp Psychol* 120:31-37
 582 McEachern M, Eadie J, van Vuren D (2007) Local genetic structure and relatedness in a solitary mammal,
 583 *Neotoma fuscipes*. *Behav Ecol Sociobiol* 61:1459-1469
 584 Morrogh-Bernard HC, Morf N, Chivers DJ, Krützen M (2011) Dispersal patterns of orang-utans (*Pongo spp.*) in
 585 a Bornean peat-swamp forest. *Int J Primatol* 32:362-376
 586 Moyer MA, McCown JW, Eason TH, Oli MK (2006) Does genetic relatedness influence space use pattern? A
 587 test on Florida black bears. *J Mammal* 87:255-261
 588 Nater A, Nietlisbach P, Arora N, van Schaik CP, van Noordwijk MA, Willems EP, Singleton I, Wich SA,
 589 Goossens B, Warren KS, Verschoor EJ, Perwitasari-Farajallah D, Pamungkas J, Krützen M (2011) Sex-
 590 biased dispersal and volcanic activities shaped phylogeographic patterns of extant orangutans (genus:
 591 *Pongo*). *Mol Biol Evol* 28:2275-2288

592 Nietlisbach P (2009) Male specific markers in orangutans (*Pongo spp.*): Dispersal and phylogeny. MSc Thesis,
 593 University of Zurich
 594 Nunes S, Muecke E-M, Sanchez Z, Hoffmeier RR, Lancaster LT (2004) Play behavior and motor development
 595 in juvenile Belding's ground squirrels (*Spermophilus beldingi*). Behav Ecol Sociobiol 56:97-105
 596 Pellis SM (2010) Conservative motor systems, behavioral modulation and neural plasticity. Behav Brain Res
 597 214:25-29
 598 Pusey A, Williams J, Goodall J (1997) The influence of dominance rank on the reproductive success of female
 599 chimpanzees. Science 277:828-831
 600 Pusey AE, Packer C (1987) Dispersal and philopatry. In: Smuts BB, Cheney DL, Seyfarth RM, Wrangham RW,
 601 Struhsaker TT (eds) Primate Societies, Chicago University Press, Chicago, pp 250-266
 602 Ratnayake S, Tuskan GA, Pelton MR (2002) Genetic relatedness and female spatial organization in a solitary
 603 carnivore, the raccoon, *Procyon lotor*. Mol Ecol 11:1115-1124
 604 Robbins A, Stoinski T, Fawcett K, and Robbins M (2009) Socioecological influences on the dispersal of female
 605 mountain gorillas—evidence of a second folivore paradox. Behav Ecol Sociobiol 63: 477-489
 606 Rodgers AR, Carr AP, Beyer HL, Smith L, Kie JG (2007) HRT: Home Range Tools for ArcGIS, Centre for
 607 Northern Forest Ecosystem Research, Ont Min Nat Res, Thunder Bay, Ontario
 608 Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM, Seyfarth RM, Cheney
 609 DL. (2009) The benefits of social capital: close social bonds among female baboons enhance offspring
 610 survival. Proc R Soc B 276:3099-3104
 611 Silk JB, Beehner JC, Bergman TJ, Crockford C, Engh AL, Moscovice LR, Wittig RM, Seyfarth RM, Cheney
 612 DL. (2010) Strong and consistent social bonds enhance the longevity of female baboons. Curr Biol
 613 20:1359-1361
 614 Singleton IS, Knott CD, Morrogh-Bernard HC, Wich SA, van Schaik CP (2009) Ranging behavior of orangutan
 615 females and social organization. In: Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP, (eds)
 616 Orangutans Geographic Variation in Behavioral Ecology and Conservation. Oxford University Press,
 617 Oxford, pp 205-213
 618 Singleton IS, van Schaik CP (2001) Orangutan home range size and its determinants in a Sumatran swamp
 619 forest. International Journal of Primatology 22:877-911
 620 Singleton IS, van Schaik CP (2002) The social organisation of a population of Sumatran orang-utans. Folia
 621 Primatol 73:1-20
 622 Smith JLD (1993) The role of dispersal in structuring the Chitwan Tiger Population. Behaviour 124:165-195
 623 Špinko M, Newberry RC, Bekoff M (2001) Mammalian play: Training for the unexpected. Q Rev Biol 76:141-
 624 168
 625 Støen O-G, Bellemain E, Sæbø S, Swenson J (2005) Kin-related spatial structure in brown bears *Ursus arctos*.
 626 Behav Ecol Sociobiol 59:191-197
 627 Townsend SW, Slocum KE, Emery Thompson M, Zuberbühler K (2007) Female-led infanticide in wild
 628 chimpanzees. Curr Biol 17:R355-R356
 629 Utami Atmoko SS, Singleton IS, van Noordwijk MA, van Schaik CP, Setia TM (2009) Male-male relationships
 630 in orangutans. In: Wich SA, Utami Atmoko SS, Setia TM, van Schaik CP (eds), Orangutans
 631 Geographic Variation in Behavioral Ecology and Conservation. Oxford University Press, Oxford, pp
 632 225-234

633 Utami SS, Goossens B, Bruford MW, de Ruiter JR, van Hooff JARAM (2002) Male bimaturism and
 634 reproductive success in Sumatran orang-utans. Behav Ecol 13:643-652
 635 van Noordwijk MA, Sauren SEB, Nuzuar, Abulani A, Morrogh-Bernard HC, Utami Atmoko SS, van Schaik CP
 636 (2009) Development of independence Sumatran and Bornean orangutans compared. In: Wich SA,
 637 Utami Atmoko SS, Mitra Setia T, van Schaik CP (eds), Orangutans: Geographic Variation in
 638 Behavioral Ecology and Conservation. Oxford University Press, Oxford, pp 189-203
 639 van Schaik CP (1999) The socioecology of fission-fusion sociality in orangutans. Primates 40:73-90
 640 van Schaik CP, van Hooff JARAM (1996) Towards an understanding of the orangutan's social system. In:
 641 McGrew WC, Marchant LF, Nishida T (eds) Great Ape Societies. Cambridge University Press,
 642 Cambridge, pp 3-15
 643 van Schaik CP, Wich SA, Utami SS, Odom K (2005) A simple alternative to line transects of nests for
 644 estimating orangutan densities. Primates 46:249-254
 645 van Woerden JT, Pettersson J (2007) The effect of reproductive state on wild female orangutan ranging behavior
 646 (*Pongo pygmaeus wurmbii*). 2nd Congr Europ Fed Primatol, Prague. p 71-72 (abstract).
 647 Wakefield ML (2008) Grouping patterns and competition among female *Pan troglodytes schweinfurthii* at
 648 Ngogo, Kibale National Park, Uganda. Int J Primatol 29:907-927
 649 Wartmann F, Purves R, van Schaik C (2010) Modelling ranging behaviour of female orang-utans: a case study in
 650 Tuanan, Central Kalimantan, Indonesia. Primates 51: 119-130
 651 Waser PM, Jones WT (1983) Natal philopatry among solitary mammals. Q Rev Biol 58: 355-390
 652 Watts P (1994) Social relationships of immigrant and resident female mountain gorillas, II:
 653 relatedness, residence, and relationships between females. Am J Primatol 32: 13-30
 654 Watts DP (2000) Causes and consequences of variation in male mountain gorilla life histories and group
 655 membership. In: Kappeler PM (ed) Primate males causes and consequences of variation in group
 656 composition. Cambridge University Press, Cambridge, pp 169-179
 657 Wich SA, Utami-Atmoko SS, Mitra Setia T, Rijksen HD, Schürmann C, van Hooff JARAM, van Schaik CP
 658 (2004) Life history of wild Sumatran orangutans (*Pongo abelii*). J Hum Evol 47:385-398
 659 Wild KD (2010) Female chimpanzees in Kanyawara community form social bonds and utilize valuable
 660 relationships despite female dispersal and male-bonded philopatry. Am J Phys Anthropol Suppl 50:243
 661 Wolff JO, Peterson JA (1998) An offspring-defense hypothesis for territoriality in female mammals. Ethol Ecol
 662 Evol 10:227-239
 663 Worton BJ (1989) Kernel methods for estimating the utilization distribution in home-range studies. Ecology
 664 70:164-168
 665

Figure legends

Figure 1

Map of the overlapping home range areas (95% use) of 7 focal females calculated for 2008. Females Jinak, Juni, Kerry and Mindy are members of the same matriline; Desy is the daughter of Inul, and for Talia, and Sidony no matrilineal adult female relatives are known. Also indicated are the parts of the home ranges of the additional females Wilma and Tina (probably maternal relatives, but not mother daughter) and Pinky for as far as they are known in the study area. The research station is indicated with a flag. (The area to the west of the study area is burnt and severely degraded habitat).

Figure 2

Comparison between maternally related females dyads and unrelated dyads in a) their home range (95% use area) and core range overlap (50% use area), b) encounter frequency (% of days with an encounter per dyad) and percentage of total focal time spent in association and c) encounter frequency and association overlap controlled for home range overlap. Unrelated dyads were only included if their home range overlap was at least 32%, the lowest overlap among related females. Significant differences between adjacent columns are indicated with *** $P < 0.0001$.

Figure 3

Comparison of social behavior during associations between related females, associations between related females lasting < 3 hours and associations between unrelated females (all lasting < 3 hr): the percentage of female-female associations with at least some social play among immatures, feeding tolerance among females (i.e. both feeding within 10m), prolongation of the association overnight, agonistic interaction between the females. Significant differences between adjacent columns are indicated with ** $P < 0.001$.

Figure 1




Figure 2





Figure 3



1 Supplemental Table 1

2 Overview of observation periods included for each parous female in the study area.

female	haplotype	2003	2004	2005	2006	2007	2008	2009	2010	Remarks
Jinak	B	x	x	x	x	x	x			2009-2010 only short follows
Juni	B				x	x	x	x	x	first offspring born early 2006
Kerry	B	x	x	x	x	x	x	x	x	
Mindy	B	x	x	x	x	x	x	x	x	
Sidony	A						x	x	x	ranges in extended study area
Sumi	A	x	x	x	x					died August 2006
Talia	A					x	x			
Desy	C			x	x	x	x	x		<2005 unhabituated
Inul	C									insufficiently habituated

3

4 Supplemental Table 2

5 Tukey's Honestly Significant Difference tests for different kinds of dyads in the duration of their associations: dyads of
6 related females compared to dyads of unrelated females, unflanged males, males of different morph or flanged males.

Bootstrap for Multiple Comparisons

Dyad type (I)	Dyad type (J)	Mean Δ_{IJ}	Sig.	95% bootstrap CI	
				Lower	Upper
Related female	Unrelated female	176.862	.000	146.975	208.255
	Unflanged male	126.318	.013	65.913	182.717
	Mixed-morph male	202.972	.000	164.150	239.430
	Flanged male	222.073	.000	193.955	251.599
Unrelated female	Related female	-176.862	.000	-208.255	-146.975
	Unflanged male	-50.545	.809	-107.402	1.274
	Mixed-morph male	26.110	.984	-5.846	53.104
	Flanged male	45.210	.836	27.641	62.468
Unflanged male	Related female	-126.318	.013	-182.717	-65.913
	Unrelated female	50.545	.809	-1.274	107.402
	Mixed-morph male	76.655	.641	20.548	136.131
	Flanged male	95.755	.327	45.916	151.600
Mixed-morph male	Related female	-202.972	.000	-239.430	-164.150
	Unrelated female	-26.110	.984	-53.104	5.846
	Unflanged male	-76.655	.641	-136.131	-20.548
	Flanged male	19.101	.996	-5.325	49.250
Flanged male	Related female	-222.073	.000	-251.599	-193.955
	Unrelated female	-45.210	.836	-62.468	-27.641
	Unflanged male	-95.755	.327	-151.600	-45.916
	Mixed-morph male	-19.101	.996	-49.250	5.325

7

8


9

10 Supplemental Figure 1

11 Comparison of % of association between different kinds of dyads during which there was social play among the
12 immatures and/or adults in association, feeding tolerances among the adults, or where the association was
13 prolonged into the next day. Abbreviations: rel. adf.: maternally related adult females with (or without) their
14 dependent offspring; rel adf <3: same but association lasting <3 hours; unrel adf : maternally unrelated females
15 with (or without) their dependent offspring; ufm-ufm: association between unflanged males; ufm-flm:
16 association between unflanged and flanged male: flm-flm: association between flanged males.


17

% Associations with Social Play




18

% Assoc. with Feeding tolerance



19

% Assoc. prolonged overnight



20